arXiv:1411.2544v1 [math.CO] 10 Nov 2014

# Randić energy of specific graphs

Saeid Alikhani[1] and Nima Ghanbari

*Department of Mathematics, Yazd University*
*89195-741, Yazd, Iran*

**Abstract**

Let $G$ be a simple graph with vertex set $V(G) = \{v_1, v_2, \ldots, v_n\}$. The Randić matrix of $G$, denoted by $R(G)$, is defined as the $n \times n$ matrix whose $(i, j)$-entry is $(d_i d_j)^{\frac{-1}{2}}$ if $v_i$ and $v_j$ are adjacent and 0 for another cases. Let the eigenvalues of the Randić matrix $R(G)$ be $\rho_1 \geq \rho_2 \geq \ldots \geq \rho_n$ which are the roots of the Randić characteristic polynomial $\prod_{i=1}^{n}(\rho - \rho_i)$. The Randić energy $RE$ of $G$ is the sum of absolute values of the eigenvalues of $R(G)$. In this paper we compute the Randić characteristic polynomial and the Randić energy for specific graphs $G$.

**Mathematics Subject Classification:** 15A18, 05C50.
**Keywords:** Randić matrix; Randić energy; Randicć characteristic polynomial; eigenvalues.

## 1 Introduction

In this paper we are concerned with simple finite graphs, without directed, multiple, or weighted edges, and without self-loops. Let $G$ be such a graph, with vertex set $V(G) = \{v_1, v_2, \ldots, v_n\}$. If two vertices $v_i$ and $v_j$ of $G$ are adjacent, then we use the notation $v_i \sim v_j$. For $v_i \in V(G)$, the degree of the vertex $v_i$, denoted by $d_i$, is the number of the vertices adjacent to $v_i$.

Let $A(G)$ be adjacency matrix of $G$ and $\lambda_1, \lambda_2, \ldots, \lambda_n$ its eigenvalues. These are said to be be the eigenvalues of the graph $G$ and to form its spectrum [4]. The energy $E(G)$ of the graph $G$ is defined as the sum of the absolute values of its eigenvalues

$$E(G) = \sum_{i=1}^{n} |\lambda_i|.$$

Details and more information on graph energy can be found in [6, 7, 9, 10].

[1]Corresponding author. E-mail: alikhani@yazd.ac.ir

In 1975 Milan Randić invented a molecular structure descriptor defined as [11]

$$R(G) = \sum_{v_i \sim v_j} \frac{1}{\sqrt{d_i d_j}}.$$

The Randić-index-concept suggests that it is purposeful to associate to the graph $G$ a symmetric square matrix $R(G)$. The Randić matrix $R(G) = (r_{ij})_{n\times n}$ is defined as [2,3,8]

$$r_{ij} = \begin{cases} \frac{1}{\sqrt{d_i d_j}} & \text{if } v_i \sim v_j \\ 0 & \text{otherwise.} \end{cases}$$

Denote the eigenvalues of the Randić matrix $R(G)$ by $\rho_1, \rho_2, \ldots, \rho_n$ and label them in non-increasing order. Similar to characteristic polynomial of a matrix, we consider the Randić characteristic polynomial of $R(G)$ (or a graph $G$), as $det(\rho I - R(G))$ which is equal to $\prod_{i=1}^{n}(\rho - \rho_i)$. The Randić energy [2,3,8] of $G$ is defined as

$$RE(G) = \sum_{i=1}^{n} |\rho_i|.$$

For several lower and upper bounds on Randić energy, see [2,3,8].

In Section 2, we obtain the Randić characteristic polynomial and energy of specific graphs. As a result, we show that for every natural number $m \geq 2$, there exists a graph $G$ such that $RE(G) = m$. In Section 3, we find Randić energy of specific graphs with one edge deleted.

As usual we denote by $J$ a matrix all of whose entries are equal to 1.

## 2 Randić characteristic polynomial and Randić energy of specific graphs

In this section we study the Randić characteristic polynomial and the Randić energy for certain graphs. The following theorem gives a relationship between the Randić energy and energy of path $P_n$.

**Lemma 1** [8] Let $P_n$ be the path on $n$ vertices. Then

$$RE(P_n) = 2 + \frac{1}{2}E(P_{n-2}).$$

The following theorem gives the Randić energy of even cycles.

**Lemma 2** [12] Let $C_{2n}$ be the cycle on $2n$ vertices for $n \geq 2$. Then

$$RE(C_{2n}) = \frac{2sin((\lfloor \frac{n}{2} \rfloor + \frac{1}{2})\frac{\pi}{n})}{sin\frac{\pi}{2n}}.$$

Here we shall compute the Randić characteristic polynomial of paths and cycles.

**Theorem 1.** *For $n \geq 5$, the Randić characteristic polynomial of the path graph $P_n$ satisfy*

$$RP(P_n, \lambda) = (\lambda^2 - 1)(\lambda\Lambda_{n-3} - \frac{1}{4}\Lambda_{n-4}),$$

*where for every $k \geq 3$, $\Lambda_k = \lambda\Lambda_{k-1} - \frac{1}{4}\Lambda_{k-2}$ with $\Lambda_1 = \lambda$ and $\Lambda_2 = \lambda^2 - \frac{1}{4}$.*

**Proof.** For every $k \geq 3$, consider

$$B_k := \begin{pmatrix} \lambda & \frac{-1}{2} & 0 & 0 & \dots & 0 & 0 & 0 \\ \frac{-1}{2} & \lambda & \frac{-1}{2} & 0 & \dots & 0 & 0 & 0 \\ 0 & \frac{-1}{2} & \lambda & \frac{-1}{2} & \dots & 0 & 0 & 0 \\ 0 & 0 & \frac{-1}{2} & \lambda & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \lambda & \frac{-1}{2} & 0 \\ 0 & 0 & 0 & 0 & \dots & \frac{-1}{2} & \lambda & \frac{-1}{2} \\ 0 & 0 & 0 & 0 & \dots & 0 & \frac{-1}{2} & \lambda \end{pmatrix}_{k\times k},$$

and let $\Lambda_k = det(B_k)$. It is easy to see that $\Lambda_k = \lambda\Lambda_{k-1} - \frac{1}{4}\Lambda_{k-2}$.

Suppose that $RP(P_n, \lambda) = det(\lambda I - R(P_n))$. We have

$$RP(P_n, \lambda) = det \left(\begin{array}{c|ccccc|c} \lambda & \frac{-1}{\sqrt{2}} & 0 & \dots & 0 & 0 & 0 \\ \hline \frac{-1}{\sqrt{2}} & & & & & & 0 \\ 0 & & & & & & 0 \\ \vdots & & & B_{n-1} & & & \vdots \\ 0 & & & & & & 0 \\ 0 & & & & & & \frac{-1}{\sqrt{2}} \\ \hline 0 & 0 & 0 & \dots & 0 & \frac{-1}{\sqrt{2}} & \lambda \end{array}\right)_{n\times n}.$$

So

$$RP(P_n,\lambda)=\lambda det\left(\begin{array}{ccc|c} & & & 0\\ & B_{n-2} & & \vdots\\ & & & 0\\ & & & \frac{-1}{\sqrt{2}}\\ \hline 0 & \dots & 0 & \frac{-1}{\sqrt{2}} \end{array}\right)+\frac{1}{\sqrt{2}}det\left(\begin{array}{c|ccc|c} \frac{-1}{\sqrt{2}} & \frac{-1}{2} & \dots & 0 & 0\\ \hline 0 & & & & 0\\ \vdots & & B_{n-3} & & \vdots\\ 0 & & & & \frac{-1}{\sqrt{2}}\\ \hline 0 & 0 & \dots & \frac{-1}{\sqrt{2}} & \lambda \end{array}\right).$$

And so

$$RP(P_n,\lambda)=\lambda(\lambda\Lambda_{n-2}+\frac{1}{\sqrt{2}}det\left(\begin{array}{ccc|c} & & & 0\\ & B_{n-3} & & \vdots\\ & & & \frac{-1}{2}\\ \hline 0 & \dots & 0 & \frac{-1}{\sqrt{2}} \end{array}\right))-\frac{1}{2}det\left(\begin{array}{ccc|c} & & & 0\\ & B_{n-3} & & \vdots\\ & & & \frac{-1}{\sqrt{2}}\\ \hline 0 & \dots & \frac{-1}{\sqrt{2}} & \lambda \end{array}\right).$$

Therefore

$$RP(P_n,\lambda)=\lambda^2\Lambda_{n-2}-\frac{1}{2}\lambda\Lambda_{n-3}-\frac{1}{2}\lambda\Lambda_{n-3}-\frac{\sqrt{2}}{4}det\left(\begin{array}{ccc|c} & & & 0\\ & B_{n-4} & & \vdots\\ & & & \frac{-1}{2}\\ \hline 0 & \dots & 0 & \frac{-1}{\sqrt{2}} \end{array}\right),$$

and so

$$RP(P_n,\lambda)=\lambda^2\Lambda_{n-2}-\lambda\Lambda_{n-3}+\frac{1}{4}\Lambda_{n-4}=\lambda^2(\lambda\Lambda_{n-3}-\frac{1}{4}\Lambda_{n-4})-\lambda\Lambda_{n-3}+\frac{1}{4}\Lambda_{n-4}.$$

Hence

$$RP(P_n,\lambda)=(\lambda^2-1)(\lambda\Lambda_{n-3}-\frac{1}{4}\Lambda_{n-4}).\quad\square$$

**Theorem 2.** *For $n \geq 3$, the Randić characteristic polynomial of the cycle graph $C_n$ is*

$$RP(C_n, \lambda) = \lambda\Lambda_{n-1} - \frac{1}{2}\Lambda_{n-2} - (\frac{1}{2})^{n-1},$$

*where for every $k \geq 3$, $\Lambda_k = \lambda\Lambda_{k-1} - \frac{1}{4}\Lambda_{k-2}$ with $\Lambda_1 = \lambda$ and $\Lambda_2 = \lambda^2 - \frac{1}{4}$.*

**Proof.** Similar to the proof of Theorem 1, for every $k \geq 3$, we consider

$$B_k := \begin{pmatrix} \lambda & \frac{-1}{2} & 0 & 0 & \dots & 0 & 0 & 0 \\ \frac{-1}{2} & \lambda & \frac{-1}{2} & 0 & \dots & 0 & 0 & 0 \\ 0 & \frac{-1}{2} & \lambda & \frac{-1}{2} & \dots & 0 & 0 & 0 \\ 0 & 0 & \frac{-1}{2} & \lambda & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \lambda & \frac{-1}{2} & 0 \\ 0 & 0 & 0 & 0 & \dots & \frac{-1}{2} & \lambda & \frac{-1}{2} \\ 0 & 0 & 0 & 0 & \dots & 0 & \frac{-1}{2} & \lambda \end{pmatrix}_{k\times k},$$

and let $\Lambda_k = det(B_k)$. We have $\Lambda_k = \lambda\Lambda_{k-1} - \frac{1}{4}\Lambda_{k-2}$.

Suppose that $RP(C_n, \lambda) = det(\lambda I - R(C_n))$. We have

$$RP(C_n, \lambda) = det \left(\begin{array}{c|ccccc} \lambda & \frac{-1}{2} & 0 & \dots & 0 & \frac{-1}{2} \\ \hline \frac{-1}{2} & & & & & \\ 0 & & & & & \\ \vdots & & & B_{n-1} & & \\ 0 & & & & & \\ \frac{-1}{2} & & & & & \end{array}\right)_{n\times n}.$$

So

$$RP(C_n, \lambda) = \lambda\Lambda_{n-1} + \frac{1}{2}det \left(\begin{array}{c|ccc} \frac{-1}{2} & \frac{-1}{2} & \dots & 0 \\ \hline 0 & & & \\ \vdots & & B_{n-2} & \\ \frac{-1}{2} & & & \end{array}\right) + (-1)^{n+1}(\frac{-1}{2})det \left(\begin{array}{c|ccc} \frac{-1}{2} & & & \\ \vdots & & B_{n-2} & \\ 0 & & & \\ \hline \frac{-1}{2} & 0 & \dots & \frac{-1}{2} \end{array}\right).$$

And so,

$$RP(C_n,\lambda)=\lambda\Lambda_{n-1}-\tfrac{1}{4}\Lambda_{n-2}+(-1)^n(\tfrac{-1}{4})det\left(\begin{array}{ccc|c}\frac{-1}{2} & \dots & 0 & 0\\ \hline & & & 0\\ & B_{n-3} & & \vdots\\ & & & \frac{-1}{2}\end{array}\right)+$$

$$(-1)^{n+1}(\tfrac{-1}{2})(\tfrac{-1}{2}det\left(\begin{array}{c|ccc}\frac{-1}{2} & & & \\ \vdots & & B_{n-3} & \\ 0 & & & \\ \hline \frac{-1}{2} & 0 & \dots & \frac{-1}{2}\end{array}\right)+(-1)^n(\tfrac{-1}{2})\Lambda_{n-2}).$$

Therefore,

$$RP(C_n,\lambda)=\lambda\Lambda_{n-1}-\frac{1}{4}\Lambda_{n-2}+(-1)^n(\frac{-1}{2})^{n-1}(\frac{1}{2})+(-1)^{n+1}(\frac{-1}{2})^n+\frac{1}{4}(-1)^{2n+1}\Lambda_{n-2}.$$

Hence

$$RP(C_n,\lambda)=\lambda\Lambda_{n-1}-\frac{1}{2}\Lambda_{n-2}-(\frac{1}{2})^{n-1}.\quad\square$$

In [12] has shown that the Randić energy of $S_n = K_{1,n-1}$, the star on $n$ vertices and the complete bipartite graph $K_{m,n}$ is 2. Here using the Randić characteristic polynomial, we prove these results. We need the following lemma:

**Lemma 3.** [4] If $M$ is a nonsingular square matrix, then

$$det\begin{pmatrix}M & N\\ P & Q\end{pmatrix}=det(M)det(Q-PM^{-1}N).$$

**Theorem 3.** *For $n \geq 2$,*

*(i) The Randić characteristic polynomial of the star graph $S_n = K_{1,n-1}$ is*

$$RP(S_n,\lambda)=\lambda^{n-2}(\lambda^2-1).$$

*(ii) The Randić energy of $S_n$ is*

$$RE(S_n)=2.$$

**Proof.**

(i) It is easy to see that the Randić matrix of $K_{1,n-1}$ is $\frac{1}{\sqrt{n-1}}\begin{pmatrix} 0_{1\times 1} & J_{1\times n-1} \\ J_{n-1\times 1} & 0_{n-1\times n-1} \end{pmatrix}$. We have

$$det(\lambda I - R(S_n)) = det\begin{pmatrix} \lambda & \frac{-1}{\sqrt{n-1}}J_{1\times(n-1)} \\ \frac{-1}{\sqrt{n-1}}J_{(n-1)\times 1} & \lambda I_{n-1} \end{pmatrix}.$$

Using Lemma 3,

$$det(\lambda I - R(S_n)) = \lambda det(\lambda I_{n-1} - \frac{1}{\sqrt{n-1}}J_{(n-1)\times 1}\frac{1}{\lambda}\frac{1}{\sqrt{n-1}}J_{1\times(n-1)}).$$

We know that $J_{(n-1)\times 1}J_{1\times(n-1)} = J_{n-1}$. Therefore

$$det(\lambda I - R(S_n)) = \lambda det(\lambda I_{n-1} - \frac{1}{\lambda(n-1)}J_{n-1}) = \lambda^{2-n}det(\lambda^2 I_{n-1} - \frac{1}{n-1}J_{n-1}).$$

Since the eigenvalues of $J_{n-1}$ are $n-1$ (once) and 0 ($n-2$ times), the eigenvalues of $\frac{1}{n-1}J_n$ are 1 (once) and 0 ($n-2$ times). Hence

$$RP(S_n, \lambda) = \lambda^{n-2}(\lambda^2 - 1).$$

(ii) It follows from Part (i). □

**Theorem 4.** *For $n \geq 2$,*

*(i) the Randić characteristic polynomial of complete graph $K_n$ is*

$$RP(K_n, \lambda) = (\lambda - 1)(\lambda + \frac{1}{n-1})^{n-1}.$$

*(ii) the Randić energy of $K_n$ is*

$$RE(K_n) = 2.$$

**Proof.**

(i) It is easy to see that the Randić matrix of $K_n$ is $\frac{1}{n-1}(J-I)$. Therefore

$$RP(K_n,\lambda) = det(\lambda I - \frac{1}{n-1}J + \frac{1}{n-1}I) = det((\lambda + \frac{1}{n-1})I - \frac{1}{n-1}J).$$

Since the eigenvalues of $J_n$ are $n$ (once) and 0 ($n-1$ times), the eigenvalues of $\frac{1}{n-1}J_n$ are $\frac{n}{n-1}$ (once) and 0 ($n-1$ times). Hence

$$RP(K_n,\lambda) = (\lambda - 1)(\lambda + \frac{1}{n-1})^{n-1}.$$

(ii) It follows from Part (i). □

**Theorem 5.** *For natural number $m, n \neq 1$,*

*(i) The Randić characteristic polynomial of complete bipartite graph $K_{m,n}$ is*

$$RP(K_{m,n},\lambda) = \lambda^{m+n-2}(\lambda^2 - 1).$$

*(ii) The Randić energy of $K_{m,n}$ is*

$$RE(K_{m,n}) = 2.$$

**Proof.**

(i) It ie easy to see that the Randić matrix of $K_{m,n}$ is $\frac{1}{\sqrt{mn}}\begin{pmatrix} 0_{m\times m} & J_{m\times n} \\ J_{n\times m} & 0_{n\times n} \end{pmatrix}$. Using Lemma 3 we have

$$det(\lambda I - R(K_{m,n})) = det\begin{pmatrix} \lambda I_m & \frac{-1}{\sqrt{mn}}J_{m\times n} \\ \frac{-1}{\sqrt{mn}}J_{n\times m} & \lambda I_n \end{pmatrix}.$$

So

$$det(\lambda I - R(K_{m,n})) = det(\lambda I_m)det(\lambda I_n - \frac{1}{\sqrt{mn}}J_{n\times m}\frac{1}{\lambda}I_m\frac{1}{\sqrt{mn}}J_{m\times n}).$$

We know that $J_{n\times m}J_{m\times n} = mJ_n$. Therefore

$$det(\lambda I - R(K_{m,n})) = \lambda^m det(\lambda I_n - \frac{1}{\lambda n}J_n) = \lambda^{m-n}det(\lambda^2 I_n - \frac{1}{n}J_n).$$

The eigenvalues of $J_n$ are $n$ (once) and 0 ($n-1$ times). So the eigenvalues of $\frac{1}{n}J_n$ are 1 (once) and 0 ($n-1$ times). Hence

$$RP(K_{m,n},\lambda) = \lambda^{m+n-2}(\lambda^2-1).$$

(ii) It follows from Part (i). □

Let $n$ be any positive integer and $F_n$ be friendship graph with $2n+1$ vertices and $3n$ edges. In other words, the friendship graph $F_n$ is a graph that can be constructed by coalescence $n$ copies of the cycle graph $C_3$ of length 3 with a common vertex. The Friendship Theorem of Erdős, Rényi and Sós [5], states that graphs with the property that every two vertices have exactly one neighbour in common are exactly the friendship graphs. The Figure 1 shows some examples of friendship graphs. Here we shall investigate the Randić energy of friendship graphs.



Figure 1: Friendship graphs $F_2, F_3, F_4$ and $F_n$, respectively.

**Theorem 6.** *For $n \geq 2$,*

*(i) The Randić characteristic polynomial of friendship graph $F_n$ is*

$$RP(F_n,\lambda) = (\lambda^2-\frac{1}{4})^{n-1}(\lambda-1)(\lambda+\frac{1}{2})^2.$$

*(ii) The Randić energy of $F_n$ is*

$$RE(F_n) = n+1.$$

**Proof.**

(i) The Randić matrix of $F_n$ is

$$R(F_n) = \begin{pmatrix} 0 & \frac{1}{2\sqrt{n}} & \frac{1}{2\sqrt{n}} & \cdots & \frac{1}{2\sqrt{n}} & \frac{1}{2\sqrt{n}} \\ \frac{1}{2\sqrt{n}} & 0 & \frac{1}{2} & \cdots & 0 & 0 \\ \frac{1}{2\sqrt{n}} & \frac{1}{2} & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \\ \frac{1}{2\sqrt{n}} & 0 & 0 & \cdots & 0 & \frac{1}{2} \\ \frac{1}{2\sqrt{n}} & 0 & 0 & \cdots & \frac{1}{2} & 0 \end{pmatrix}_{(2n+1)\times(2n+1)}.$$

Now for computing $det(\lambda I - R(F_n))$, we consider its first row. The cofactor of the first array in this row is

$$\begin{pmatrix} \lambda & \frac{-1}{2} & \cdots & 0 & 0 \\ \frac{-1}{2} & \lambda & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \lambda & \frac{-1}{2} \\ 0 & 0 & \cdots & \frac{-1}{2} & \lambda \end{pmatrix}$$

and the cofactor of another arrays in the first row are similar to

$$\begin{pmatrix} \frac{-1}{2\sqrt{n}} & \frac{-1}{2} & \cdots & 0 & 0 \\ \frac{-1}{2\sqrt{n}} & \lambda & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{-1}{2\sqrt{n}} & 0 & \cdots & \lambda & \frac{-1}{2} \\ \frac{-1}{2\sqrt{n}} & 0 & \cdots & \frac{-1}{2} & \lambda \end{pmatrix}$$

Now, by straightforward computation we have the result.

(ii) It follows from Part (i). □

**Remark.** In [1] has shown that the energy of a graph cannot be an odd integer. Since $RE(F_n) = n + 1$ for $n \geq 2$, the Randić energy can be odd or even integer. More precisely we have:

**Corollary 1.** *For every natural number $m \geq 2$, there exists a graph $G$ such that $RE(G) = m$.*

**Proof.** If $m = 2$ then consider $K_2$ and for $m \geq 3$ we consider friendship graphs $F_{m-1}$. □

Let $n$ be any positive integer and $D_4^n$ be Dutch Windmill Graph with $3n + 1$ vertices and $4n$ edges. In other words, the graph $D_4^n$ is a graph that can be constructed by coalescence $n$ copies of the cycle graph $C_4$ of length 4 with a common vertex. The Figure 2 shows some examples of Dutch Windmill graphs. Here we shall investigate the Randić energy of Dutch Windmill graphs.




Figure 2: Dutch Windmill Graph $D_4^2, D_4^3, D_4^4$ and $D_4^n$, respectively.

**Theorem 7.** *For $n \geq 2$,*

*(i) The Randić characteristic polynomial of friendship graph $D_4^n$ is*

$$RP(D_4^n, \lambda) = \lambda^{n+1}(\lambda^2 - \frac{1}{2})^{n-1}(\lambda^2 - 1).$$

*(ii) The Randić energy of $F_n$ is*

$$RE(D_4^n) = 2 + (n-1)\sqrt{2}.$$

**Proof.**

(i) The Randić matrix of $D_4^n$ is

$$R(D_4^n) = \begin{pmatrix} 0 & \frac{1}{2\sqrt{n}} & \frac{1}{2\sqrt{n}} & 0 & \cdots & \frac{1}{2\sqrt{n}} & \frac{1}{2\sqrt{n}} & 0 \\ \frac{1}{2\sqrt{n}} & 0 & 0 & \frac{1}{2} & \ldots & 0 & 0 & 0 \\ \frac{1}{2\sqrt{n}} & 0 & 0 & \frac{1}{2} & \ldots & 0 & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 & \ldots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \\ \frac{1}{2\sqrt{n}} & 0 & 0 & 0 & \ldots & 0 & 0 & \frac{1}{2} \\ \frac{1}{2\sqrt{n}} & 0 & 0 & 0 & \ldots & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 0 & \ldots & \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix}_{(3n+1)\times(3n+1)}.$$

Let $A = \begin{pmatrix} \lambda & 0 & \frac{-1}{2} \\ 0 & \lambda & \frac{-1}{2} \\ \frac{-1}{2} & \frac{-1}{2} & \lambda \end{pmatrix}$, $B = \begin{pmatrix} \frac{-1}{2\sqrt{n}} & 0 & 0 \\ \frac{-1}{2\sqrt{n}} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and $C = \begin{pmatrix} \frac{-1}{2\sqrt{n}} & 0 & \frac{-1}{2} \\ \frac{-1}{2\sqrt{n}} & \lambda & \frac{-1}{2} \\ 0 & \frac{-1}{2} & \lambda \end{pmatrix}$.

Then

$$det(\lambda I - R(D_4^n)) = \lambda(det(A))^n + 2n.\frac{1}{2\sqrt{n}} det \begin{pmatrix} C & 0 & 0 & \ldots & 0 \\ B & A & 0 & \ldots & 0 \\ B & 0 & A & \ldots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ B & 0 & 0 & \ldots & A \end{pmatrix}_{3n\times 3n}.$$

Now, by the straightforward computation we have the result.

(ii) It follows from Part (i). □

## 3 Randić energy of specific graphs with one edge deleted

In this section we obtain the randić energy for certain graphs with one edge deleted. We need the following lemma:

**Lemma 4.** *Let $G = G_1 \cup G_2 \cup \ldots \cup G_m$. Then*

$$RE(G) = Re(G_1) + RE(G_2) + \ldots + RE(G_m).$$

Here we state the following easy results:

**Lemma 5.**

*(i) If $e \in E(P_n)$, then $RE(P_n - e) = RE(P_r) + RE(P_s)$, where $r + s = n$.*

*(ii) If $e \in E(C_n)$, $(n \geq 3)$, then $RE(C_n - e) = RE(P_n)$.*

*(iii) Let $S_n$ be the star on $n$ vertices and $e \in E(S_n)$. Then for any $n \geq 3$,*

$$RE(S_n - e) = RE(S_{n-1}) = 2.$$

**Theorem 8.** *For $n \geq 2$,*

*(i) Let $e$ be an edge of complete graph $K_n$. The Randić characteristic polynomial of $K_n - e$ is*

$$RP(K_n - e, \lambda) = \lambda(\lambda - 1)(\lambda + \frac{2}{n-1})(\lambda + \frac{1}{n-1})^{n-3}.$$

*(ii) The Randić energy of $K_n - e$ is*

$$RE(K_n - e) = 2.$$

**Proof.**

(i) The Randić matrix of $K_n - e$ is

$$\begin{pmatrix} 0_{2\times 2} & \frac{1}{\sqrt{(n-1)(n-2)}} J_{2\times(n-2)} \\ \frac{1}{\sqrt{(n-1)(n-2)}} J_{(n-2)\times 2} & \frac{1}{n-1}(J - I)_{n-2} \end{pmatrix}.$$

Therefore

$$det(\lambda I - R(K_n - e)) = det \begin{pmatrix} \lambda I_2 & \frac{-1}{\sqrt{(n-1)(n-2)}} J_{2\times(n-2)} \\ \frac{-1}{\sqrt{(n-1)(n-2)}} J_{(n-2)\times 2} & \frac{\lambda}{n-1}(J - I)_{n-2} \end{pmatrix}.$$

Similar to the proof of Theorem 5(i) we have the result.

(ii) It follows from Part (i). □

**Theorem 9.** *For $m, n \neq 1$,*

*(i) The Randić characteristic polynomial of complete bipartite graph $K_{m,n} - e$ is*

$$RP(K_{m,n} - e, \lambda) = \lambda^{m+n-4}(\lambda^2 - 1)(\lambda^2 - \frac{1}{mn}).$$

*(ii) The Randić energy of $K_{m,n} - e$ is*

$$RE(K_{m,n} - e) = 2 + \frac{2}{\sqrt{mn}}.$$

**Proof.**

(i) The Randić matrix of $K_{m,n} - e$ is

$$\begin{pmatrix} 0_{m\times m} & A \\ A^t & 0_{n\times n} \end{pmatrix},$$

where $A = \begin{pmatrix} \frac{1}{\sqrt{mn}} J_{(m-1)\times(n-1)} & \frac{1}{\sqrt{n(m-1)}} J_{(m-1)\times 1} \\ \frac{1}{\sqrt{m(n-1)}} J_{1\times(n-1)} & 0_{1\times 1} \end{pmatrix}$.So

$$det(\lambda I - R(K_{m,n} - e)) = \lambda I_m det(\lambda I_n - A^t \frac{1}{\lambda} I_m A).$$

Similar to the proof of Theorem 5(i) we have the result.

(ii) It follows from Part (i). □

## References


[1] R. B. Bapat, S. Pati, Energy of a graph is never an odd integer, *Bull. Kerala Math. Assoc.*, 1 (2004), 129–132.

[2] Ş. B. Bozkurt, A. D. Güngör, I. Gutman, A. S. Çevik, Randić matrix and Randić energy, *MATCH Commum. Math. Comput. Chem.* 64 (2010) 239–250.

[3] Ş. B. Bozkurt, A. D. Güngör, I. Gutman, Randić spectral radius and Randić energy, *MATCH Commum. Math. Comput. Chem.* 64 (2010) 321–334.

[4] D. Cvetković, M. Doob, H. sachs, *Spectra of graphs* - Theory and Aplication, Academic Press, New York, 1980.

[5] P. Erdös, A. Rényi, V.T. Sós, On a problem of graph theory, *Studia Sci. Math. Hungar.*, 1, 215–235 (1966).

[6] I. Gutman, The energy of a graph: Old and new results, in: A. Betten, A.Kohnert, R. Laue, A. Wassermannn (Eds.), *Algebraic Combinatorics and Applications*, Springer-Verlag, Berlin, 2001, pp. 196–211.

[7] I. Gutman, Topology and stability of conjugated hydrocarbons. The dependence of total $\pi$-electron energy on molecular topology, *J. Serb. Chem. Soc.* 70 (2005) 441–456.

[8] I. Gutman, B. Furtula, Ş. B. Bozkurt, On Randić energy, *Linear Algebra Appl.*, 442 (2014) 50–57.

[9] I. Gutman, X. Li, J. Zhang, Graph energy, in: M. Dehmer, F. Emmert-Streib (Eds.), *Analysis of Complex Networks. From Biology to Linguistics*, Wiley-VCH, Weinheim, 2009, pp. 145–174.

[10] S. Majstorović, A. Klobucar, I. Gutman, Selected topics from the theory of graph energy: hypoenergetic graphs, in: D. Cvetković, I. Gutman (Eds.), *Applications of Graph Spectra*, Math. Inst., Belgrade, 2009, pp. 65–105.

[11] M. Randić, On characterization of molecular branching, *J. Amer. Chem. Soc.* 97 (1975) 6609–6615.

[12] O. Rojo, L. Medina, Construction of bipartite graphs having the same Randić energy, *MATCH Commun. Math. Comput. Chem.* 68 (2012) 805–814.